

FREQUENCY: 2

ISSN: 1817 - 3055

WORLD JOURNAL OF MEDICAL SCIENCES

W J M S



Molecular Identification of Some *Schistosoma mansoni* Isolates in Saudi Arabia

Manal B. Jamjoom

Department of Medical Parasitology, Faculty of Medicine,
King Abdul-Aziz University, Jeddah, KSA

Abstract: In this paper, nine geographical isolates of *Schistosoma mansoni* (three from Egypt, three from Saudi Arabia and 3 from Puerto Rico) were studied at the genotype level by RAPD analysis with two arbitrary primers. The genetic distance was measured by the percentage of unshared bands. The RAPD results showed that the Egyptian strains were closely related to Saudi strains but Puerto Rico strains clustered in different group. These results demonstrated the usefulness of RAPD for displaying the differences of inter- and intra-species of *Schistosoma*. It also suggested that genetic diversity among different geographical strains of *Schistosoma mansoni* from different localities had occurred.

Key words: *Schistosoma mansoni* · RAPD · Egypt · genetic variation

INTRODUCTION

Schistosomiasis is caused by infection with blood flukes of the genus *Schistosoma*, of which three species (*S. mansoni*, *S. haematobium* and *S. japonicum*) are the main causative agent of the disease in man [1]. This chronic helminthic infection affects more than 200 million people throughout the world and 600 million live in endemic areas where they are at risk of infection [2]. This wide distribution of the disease makes the study of genomic variability extremely important.

There are well known two distinct geographical strains of *Schistosoma mansoni*, the Puerto Rican and the Egyptian strains [3-5]. However, a new geographical strain namely, Saudi Arabian strain has been reported based on differences with the Egyptian strain shown by scanning electron microscopy [6, 7].

Early studies of population structure in schistosomes used isozymes to look for genetic differences between populations of *S. mansoni*: those that are transmitted through rodents and populations that infected humans [8, 9]. A substantial amount of phenotypic variation has been observed in schistosome species, it has been more difficult to quantify the degree of genetic differentiation among these same populations [10]. More rigorous studies of population structure and genetic subdivision are needed to clarify our understanding of schistosome epidemiology [11]. Ideally, such studies would employ

genetic markers that are highly polymorphic and inherited in a Mendelian codominant fashion. The data available from isoenzyme analysis of laboratory strains of schistosomes from Africa, Asia, the Caribbean and Brazil [12] indicate that there is relatively restricted diversity in these species and strains with no distinct genetic constitutions.

Several studies have shown that the RAPD technique can be applied successfully to helminths [13-19] for estimating differentiation among parasite populations [20]. Recently, there have been important advances in the application of recombinant DNA techniques. A number of DNA-based methods have been developed to investigate genetic diversity, to differentiate strains and species and to analyze phylogenies of schistosomes. Dias-Neto *et al.* [13] have demonstrated that RAPD (random amplified polymorphic DNA markers) allows the identification of strains of *S. mansoni* and species of *Schistosoma*. They reported that, random amplified polymorphic DNA (RAPD) showed very discrete variability in different strains of *S. mansoni* adult worms, however, all of them were derived from laboratory maintained populations [18]. Pillay *et al.* [21] have also demonstrated intraspecific DNA polymorphisms among the isolates of *S. mansoni*.

The aim of this investigation was to confirm the differences between the closely similar isolates of

Corresponding Author: Dr. Manal B. Jamjoom, King AbdulAziz University, Department of Medical Parasitology / Female Section, P.O. Box 42806, Jeddah 21551, Saudi Arabia

Schistosoma mansoni isolated from different localities (Saudi Arabia, Egypt and Puerto Rico) in which the morphological and physiological characters were not enough for obvious separation.

MATERIALS AND METHODS

Parasite isolates: Nine isolates of repeated twice *S. mansoni* were obtained from different places: SMS1,2,3 (from Saudi Arabia); SME1,2,3 (from Egypt) and SMP1,2,3 (from Puerto Rico).

Isolation of genomic DNA: DNA was extracted from individual adult worms as follows: frozen worms were individually homogenized with a pestle in 100 μ L of 10 mM Tris-HCl pH8; 1 mM EDTA, 10 mM NaCl and 70 mM sucrose (extraction buffer). The pestle was rinsed in an Eppendorf tube with 10% sodium dodecyl sulphate and 12 μ L of proteinase (10 mg mL⁻¹). The homogenate was incubated at 57°C for 2 h, then extracted once with an equal volume of phenol and once with chloroform. The solution was adjusted to 0.3 M with sodium acetate and precipitated at 20°C overnight with 2 volumes of absolute ethanol. DNA was pelleted the following day by centrifugation (15000 g, 20 min, 4°C), then rinsed with ethanol 70%, dried and finally resuspended in 100 μ L of TE (10 mM Tris-HCl, pH8 and 1 mM EDTA). This extraction protocol yielded sufficient DNA for approximately 12 RAPD reactions worm [22].

RAPD - PCR amplification: Two arbitrary primers A7 and A8 (5'-GAAACAAATG-3' and 5'-GTGACGTAGG-3', respectively) were used as described [23]. The random amplification procedure was performed essentially as described [24] PCR reaction mixture was subjected to electrophoresis in a 1% agar gel and photographed.

Data analysis: Computer analysis of RAPD patterns was performed as given by Halmschlager *et al.* [25]. Basically, the formation obtained from agarose gel electrophoresis was digitalized by hand to a two - discrete - character - matrix (0 and 1 for absence and presence of RAPD bands). Dendrogram was calculated by using the Jukes - Cantor option in the DNADIST program and application of the FITCH program to the computed distance matrix (PHYLIP package [26]). For running DNADIST, the two discrete characters of 0 and 1 had to be converted to Guanine and Thymine in the RAPD data matrix. Complete alignment of data was performed with CLUSTALX software and then the cluster analysis will be ready by using Treecon programme [27].

RESULTS AND DISCUSSION

Nine isolates of *S. mansoni* were used in this study. The two primers (5'-GAAACAAATG-3' and 5'-GTGACGTAGG-3') used in this study, generated a considerable number of amplification products for comparison. A different DNA banding pattern was present in almost every isolates. Comparison of each profile for each of the primers was based on the presence (1) versus absence (0) of RAPD amplimers that migrated to the same position in the gel. Bands of the same size obtained by the same primer were scored as identical, but only bands repeatable in at least two experiments with the same primer at different times were evaluated. All two primers revealed high similarities between each type of isolates (Figs. 1 & 2). Anou *et al.* [28] reported that isolates of *Schistosoma japonicum* showed identical RAPD patterns by using seven primers.

The combined data from all isolates of *S. mansoni* by using two different primers were analyzed to produce a dendrogram (Fig. 3). According to dendrogram constructed from RAPD data, the isolates of *Schistosoma mansoni* isolates split into two RAPD groups (RAPD I and RAPD II). RAPD I cluster divided into two subgroups RAPD I A and RAPD I B. RAPD I A included *S. mansoni* isolates from Saudi Arabia (SMS) and RAPD I B included *S. mansoni* isolates from Egypt (SME). This result indicates that those two groups of isolates related and slightly far from the Puerto Rico isolates which clustered in RAPD II. Dendrogram revealed a correlation between clusters and geographical origin of isolates. Those results revealed a clear and significant genetic differentiation among the three local populations of adult schistosomes. These came in agreement with findings of Sire *et al.* [22]. Differences in snail infectivity [29, 30] and drug susceptibility [31-33] were also reported. The study of ribosomal DNA (rDNA) has shown that schistosomes exhibit both interspecific and intraspecific differences. It has been suggested that the variation in the major tandemly repeated copies of the gene can be used for species identification [34-37], whereas low copy numbers variants that exhibit intraspecific polymorphism can be used for strain identification [34, 35, 38].

This study has shown that there is considerable genotypic and phenotypic variability among *S. mansoni* isolates obtained from different geographic regions. When applied to the study of Schistosomes, the RAPD markers method has proved useful in analyzing the different problems associated with their genetic diversity. Its main additional advantages over the other techniques




Fig. 1: RAPD fragments generated by the primer A7 (5'-GAAACAAATG-3') for nine isolates of *Schistosoma mansoni* from Saudi Arabia (SMS), Egypt (SME) and from Puerto Rico (SMP)




Fig. 2: RAPD fragments generated by the primer A8 (5'-GTGACGTAGG-3') for nine isolates of *Schistosoma mansoni* from Saudi Arabia (SMS), Egypt (SME) and from Puerto Rico (SMP)




Fig. 3: The dendrogram showing the relationships of different *Schistosoma* isolates based in two different primers results

are speed, reproducibility, technical simpleness, high resolution and considerable reduction of the amount of DNA for routine analysis. Using this technique, markers which can be used to discriminate between species or strains were revealed quickly and without the need for sequence information or radiolabelling.

In conclusion, RAPD markers are highly resolving and helpful tool for investigation of genetic variation within the genus *Schistosoma* and the species *S. mansoni*. They provide a single technology that can be used to rapidly distinguish species and strains. Compared with other techniques of biochemistry and molecular biology, we can understand this endemic disease better.

REFERENCES

1. Rollinson, D. and V.R. Southgate, 1987. The genus *Schistosoma*: a taxonomic appraisal. In D. Rollinson and A.J. Simpson (Eds.). *The Biology of Schistosomes: from Genes to Latrines*. Academic Press, London, pp: 1-49.
2. Mott, K.E., 1987. Schistosomiasis control. In D. Rollinson and A.J. Simpson (Eds.), *The Biology of Schistosomes: from Genes to Latrines*. Academic Press, London, pp: 425-431.
3. Taylor, M.G. and G.S. Nelson, 1971. A comparison of susceptibility to niridazole of two geographical strains of *Schistosoma mansoni* in mice with a note on the susceptibility of *S. mattheei*. *Trans. Royal Soc. Trop. Med. & Hygiene*, 65: 169-174.
4. Voge, M. and N.S. Mansour, 1980. An unusual structural feature of the Egyptian strain of *Schistosoma mansoni*. *J. Parasitol.*, 66: 862-863.
5. Saoud, M.F., 1966. The infectivity and pathogenicity of geographical strains of *Schistosoma mansoni*. *Trans. Royal Soc. Trop. Med. & Hyg.*, 60: 583-600.
6. Shalaby, I.M., A.A. Banaja and A.M. Ghandour, 1991. Scanning electron microscopy of the tegumental surface of in vivo treated *Schistosoma mansoni* (Saudi Arabian geographical strain) with oxamniquine and praziquantel. *J. Egypt. Soc. Parasitol.*, 21: 787-810.
7. Shalaby, I.M., A.A. Banaja and A.M. Ghandour, 1993. Comparative scanning electron microscopy study of the tegumental surface of *Schistosoma mansoni* (Egyptian and other strains) in mice. *J. Egypt. Ger. Soc. Zool.*, 12: 27-41.
8. Rollinson, D., V.R. Southgate, J. Vercruyse and P.J. Moore, 1986. *Schistosoma mansoni* from naturally infected *Rattus rattus* in Guadeloupe: identification, prevalence and enzyme polymorphism. *Parasitology*, 93: 39-53.
9. Sene, M., 1997. Comparison of human and murine isolates of *Schistosoma mansoni* from Richard-Toll, Senegal, by isoelectric focusing. *J. Helminthol.*, 71: 175-181.
10. Minchella, D.J., R.E. Sorensen, J. Curtis and A.A. Bieberich, 1997. Molecular biology of trematodes: advances and applications. In: *Advances in Trematode Biology* (Eds., Fried, B. and T.K. Graczyk). CRC Press, Boca Raton, FL., pp: 405-446.

11. Curtis, J. and D.J. Minchella, 2000. Schistosome population genetic structure: when clumping worms is not just splitting hairs. *Parasitology Today*, 16: 68-71.
12. Fletcher, M., P.T. LoVerde and D.S. Woodruff, 1981. Genetic variation in *Schistosoma mansoni*: enzyme polymorphisms in population from Africa, Southwest Asia, South America and the West Indies. *Am. J. Trop. Med. Hyg.*, 30: 406-421.
13. Dias Neto E., C. Pereira De Souza, D. Rollinson, N. Katz, S.D.J. Pena and A.J.G. Simpson, 1993. The random amplification of polymorphic DNA allows the identification of strains and species of schistosomes. *Molecular and Biochemical Parasitology*, 57: 83-88.
14. Barral, V., S. Morand, J.P. Pointier, A. Theron, 1996. Distribution of schistosome genetic diversity within naturally infected *Rattus rattus* detected by RAPD markers. *Parasitology*, 113: 511-517.
15. Gasser, R.B., Q. Bao-Zhen, P. Nansen, M.V. Johansen and H. Bogh, 1996. Use of RAPD for the detection of genetic variation in the human blood fluke, *Schistosoma japonicum*, from mainland China. *Molecular and Cellular Probes*, 10: 353-358.
16. Dabo, A., P. Durand, S. Morand, J. Languard, D. Imbertestablet, O. Doumbo and J. Jourdane, 1997. Dispersion and genetic diversity of *Schistosoma haematobium* within its Bulinid intermediate hosts in Mali. *Acta Tropica*, 66: 15-26.
17. Fisher, M.C. and M.E. Viney, 1998. The population genetic structure of the facultatively sexual parasitic nematode *Strongyloides ratti* in wild rats. *Proceedings of the Royal Society of London, B* 265, 703-709. *Zoology* 67: 2021-2025.
18. Sire, C., P. Durand, J.P. Pointier and A. Theron, 1999. Genetic diversity and recruitment pattern of *Schistosoma mansoni* in a *Biomphalaria glabrata* snail population: a field study using random-amplified polymorphic DNA markers. *J. Parasitol.*, 85: 436-441.
19. Davies, C.M., J.P. Webster, O. Kruger, A. Munatsi, J. Ndamba and M.E.J. Woolhouse, 1999. Host-parasite population genetics, a cross-sectional comparison of *Bulinus globosus* and *Schistosoma haematobium*. *Parasitology*, 119: 295-302.
20. Nadler, S.A., R. Lindquist and T.J. Near, 1995. Genetic structure of midwestern *Ascaris suum* populations: a comparison of isoenzyme and RAPD markers. *J. Parasitol.*, 81: 385-394.
21. Pillay, D. and B. Pillay, 1994. Random amplified polymorphic DNA analysis shows intraspecies variation among *Schistosoma mansoni* isolates. *Med. Sci. Res.*, 22: 369-371.
22. Sire, A., P. Durand, J.P. Pointier and A. Theron, 2001. Genetic diversity of *Schistosoma mansoni* within and among individual hosts (*Rattus rattus*): infrapopulation differentiation at microspatial scale. *Intl. J. Parasitol.*, 31: 1609-1616.
23. Barral, V., P. This, D. Imbert-Establet, C. Combes and M. Delseny, 1993. Genetic variability and evolution of the *Schistosoma* genome analysed by using random amplified polymorphic DNA markers. *Molecular and Biochemical Parasitology*, 59: 211-222.
24. Williams, J.G.K., A.R. Kubelik and K.J. Livak, 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.*, 18: 6531-6535.
25. Halmschlager, E., R. Messner, T. Kowalski and H. Prillinger, 1994. Differentiation of *Ophiostoma piceae* and *Ophiostoma quercus* by morphology and RAPD-analysis. *System. Appl. Microbiol.*, 17: 554-562.
26. Felsenstein, J., 1989. PHYLIP - Phylogeny Inference Package (Version 3.2). *Cladistics*, 5: 164-166.
27. Van de Peer, Y., 1994. User manual for Treecon. Version 3.0, a software package for the constructing and drawing of evolutionary trees. University of Antwerp; Antwerp, Belgium.
28. Anou, N., X. Yanwen and F. Youren, 2002. *Schistosoma japonicum* strains: differentiation by RAPD and SSR-PCR. *Southeast Asian J. Trop. Med. Public Health*, 33: 720-724.
29. Paraense, W.L. and L.R. Correa, 1963. Variation in susceptibility of populations of *Australorbis glabratus* to a strain of *Schistosoma mansoni*. *Rev. Inst. Med. Trop. São Paulo*, 5: 15-22.
30. Richards, C.S. and P.C. Shade, 1987. The genetic variation of compatibility in *Biomphalaria glabrata* and *Schistosoma mansoni*. *J. Parasitol.*, 73: 1146-1151.
31. Katz, N., E.P. Dias, N. Araujo and C.P. Souza, 1973. Estudo de uma cepa humana de *Schistosoma mansoni* resistente a agentes esquistossomicidas. *Rev. Soc. Bras. Med. Trop.*, 7: 382-387.
32. Araujo, N., N. Katz, E. Pinto Dias and C.P. de Souza, 1980. Susceptibility to chemotherapeutic agents of strain of *Schistosoma mansoni* isolated from treated and untreated patients. *Am. J. Trop. Med. Hyg.*, 29: 890-894.

33. Dias, L.C.S., P.R. Jesus and E.R. Deberaldini, 1982. Use of praziquantel in patients with schistosomiasis mansoni previously treated with oxamniquine and/or hycanthon: resistance of *Schistosoma mansoni* to schistosomicidal agents. *Trans. R. Soc. Trop. Med. Hyg.*, 76: 652-659.
34. Simpson, A.J., J.B. Dame, F.A. Lewis and T.F. McCutchan, 1984. The arrangement of ribosomal RNA genes in *Schistosoma mansoni*. Identification of polymorphic structural variants. *Eur. J. Biochem.*, 139: 41-45.
35. McCutchan, T.F., A.J. Simpson, J.A. Mullins, A. Sher, T.E. Nash, F. Lewis and C. Richards, 1984. Differentiation of schistosomes by species, strain and sex by using cloned DNA markers. *Proc. Natl. Acad. Sci., USA.*, 81: 889-893.
36. Walker, T.K., D. Rollinson and A.J. Simpson, 1986. Differentiation of *Schistosoma haematobium* from related species using cloned ribosomal RNA probes. *Mol. Biochem. Parasitol.*, 20: 123-131.
37. Walker, T.K., A.J. Simpson and D. Rollinson, 1989. Differentiation of *Schistosoma mansoni* from *S. rodhaini* using cloned DNA probes. *Parasitology*, 98: 75-80.
38. Vieira, I.Q., R. Correa-Oliveira, N. Katz, C.P. De Souza, O.S. Carvalho, N. Araujo, A. Sher and P.J. Brindley, 1991. Genomic variability in field populations of *S. mansoni* in Brazil as detected with a ribosomal idosigene probe. *Am. J. Trop. Med. Hyg.*, 44: 69-78.